

PROTECTING TODDLERS FROM TOOTH DECAY GUIDANCE & TIPS FOR PARENTS



You can protect your toddler's teeth and prevent tooth decay by regular brushing with a fluoride toothpaste, and sensible eating and drinking habits.

Regular effective tooth brushing to remove plaque

Start brushing your baby's teeth as soon as they erupt and register him or her with a dentist. Brush twice each day – once in the morning and once at night at bedtime.

An easy way to brush your child's teeth is from behind. Comfortable and safe positions for your toddler can be on your knee, sitting on a changing mat, in a high chair or in a pram or buggy.

As more teeth come through, develop a system of cleaning them that ensures all surfaces are brushed thoroughly twice each day.

Toddlers should be encouraged to enjoy brushing their own teeth, but children should be supervised until at least seven years and always taught to 'spit don't rinse', because excessive rinsing removes the benefit of fluoride.

Fluoride toothpaste to strengthen tooth enamel

Brush teeth, twice a day with a small soft dry brush using a toothpaste containing fluoride

- **up to the age of three** use a smear of toothpaste containing 1000ppm of fluoride
- **over the age of three** use a pea sized amount of toothpaste containing 1350-1500ppm of fluoride

Take care with sweet food and drinks

Do not give sweet drinks including fruit juices and smoothies to your child. If used, dilute fruit juices because they are both sweet and acidic and can damage teeth. Serve them in a cup, never in a bottle.

Never leave a bottle of milk or a sweet drink with your toddler when you put him or her down to sleep.

Begin using a cup or free-flow beaker for drinks from about six months or so, and stop using bottles around 12 months. Never give sweet drinks in a bottle because sucking slowly on sweet drinks increases the risk of tooth decay.

Water or milk are safe drinks between meals and snacks. Give only water at night to toddlers.

Limit foods and drinks containing sugar to four times a day, e.g. at the three meals and one snack. The other snacks should be savoury snacks with no sweet food or drinks.

Do not allow grazing on food and drinks throughout the day. Only offer your toddler food at three meals and two to three snacks each day.

If your child uses a pacifier, soother or dummy make sure it is an orthodontic dummy and never dip it in anything sweet.

Nutritious snacks

Fruit	mouth-sized chunks, cubes or slices of apple, pear, satsuma, orange, banana, kiwi, melon, strawberries, sliced grapes, peaches and plums
Raw vegetable sticks or slices	carrot, pepper, celery, cucumber, sliced cherry tomato, radish, broccoli
Toast, breads, rolls, baps, bread sticks, plain buns	French bread with a small amount of butter or spread; mini savoury sandwiches with marmite, cheese, tuna, banana, salad or thin slices of meat; pita bread pockets cut into small slices or toasted with or without butter or spread e.g. olive or sunflower oil spread; breadsticks on their own or with a dip or soft cheese
Oatcakes, rice cakes, crackers, crispbreads, melba toast	plain or with butter or spread e.g. olive oil or sunflower oil spread, or use a savoury spread, such as marmite or peanut butter
Natural yogurt or fromage frais	on its own or as a dip or with fruit
Muffins, plain, potato or cheese scones, crumpets, pancakes, plain popcorn	on their own or with a small amount of butter or fat spread
Wholegrain breakfast cereals, but not cereal bars	serve with milk and fresh fruit

Where possible use a sugar-free medicine

If your child is prescribed a medicine, ensure that it is a sugar-free (SF) medicine if possible. If there is no sugar-free option give the medicine at meal times.



The healthiest start in life
from pregnancy to preschool

The Infant & Toddler Forum CIC is committed to a world where every child has the healthiest start in life

Copyright rests with the Infant & Toddler Forum CIC
Infant & Toddler Forum CIC. Registered in England and Wales. Company limited by guarantee. Company no. 12303732